

## Who can administer

May be administered by registered competent doctor or nurse/midwife

## Important information

- See **monitoring** requirements - see below
- There are numerous important **interactions** - check current BNF
- See under 'Dose' for adjustments required in **renal** impairment

## Available preparations

Rifadin 600mg vial

## Reconstitution

### Solvent provided

- 10mL solvent per 600mg vial
- Swirl vial gently to completely dissolve powder
- The solution is red in colour

### Dilute further prior to administration

## Infusion fluids

Sodium chloride 0.9% or Glucose 5%

## Methods of intravenous administration

### Intermittent intravenous infusion

- Add required dose to 500mL infusion fluid and administer over 2 to 3 hours
- Fluid restricted (use a large vein): 600mg in 100mL infusion fluid over 30 minutes (unlicensed). This solution is less stable, watch closely for precipitation <sup>(ref 1)</sup>

## Dose in adults

### Tuberculosis, in combination with other anti-tuberculous drugs

- Give 8 to 12mg/kg (to a maximum of 600mg <sup>(ref 3)</sup>) as a single daily dose
- A reduced dose (e.g. 8mg per kg daily) is recommended in elderly or frail patients

**Non-tuberculous infections** (for example: Brucellosis, Legionnaires disease, endocarditis and serious staphylococcal infections (in combination with other drugs))

- For patients over 50kg: Give 450mg twice daily, or 300mg three times daily

### Hepatic impairment

- A daily dose of 8mg/kg should not be exceeded in patients with impaired liver function

## Renal impairment <sup>(ref 2)</sup>

- If eGFR less than 10mL/min/1.73m<sup>2</sup>, use 50 to 100% of dose (to a maximum of 600mg daily)
- There is no increase in half-life in severe renal impairment for doses less than 600mg daily

## Monitoring

- Monitor baseline **LFTs, FBC and platelets**. Monitor Bilirubin, serum creatinine
- Hypersensitivity phenomena may occur (affecting platelets, vascular tissues and renal function)
- Anaphylaxis may occur, especially with intermittent therapy (e.g. two to three times weekly dosing)
- If hepatic impairment, use with caution. Use lower doses and monitor LFTs every two to four weeks during therapy

## Further information

- **Avoid extravasation** during injection; local irritation and inflammation due to extravascular infiltration of the infusion have been observed. If these occur, the infusion should be discontinued and restarted at another site
- May cause red discolouration of urine, sweat, sputum and tears

## Storage

- Store below 25<sup>0</sup>C
- The prepared infusion bag must be used within 6 hours

## References

SPC June 2022

(1) Injectable Medicines Administration Guide Medusa, downloaded 28/07/2022

(2) Renal Drug Database accessed online 28/07/2022

## Therapeutic classification

Antibiotic