

Isosorbide dinitrate Intravenous Infusion for Adults

Who can administer

Administration RESTRICTED - see [Appendix 1](#)

Important information

- See monitoring requirements below
- **Adsorbs on to PVC and PU** (polyurethane)- see under Methods of intravenous administration and also Further information
- For Y-site compatibility - [see below](#)

Available preparations

Isoket 10mg per 10ml (0.1%) ampoule

Reconstitution

Already in solution

Ampoule (10mg per 10ml)

- **Dilute further prior to administration**
- **Draw up using a 5 micron filter needle**

Infusion fluids

Sodium chloride 0.9% or Glucose 5%

Methods of intravenous administration

Continuous intravenous infusion (administer using an electronically controlled infusion device)

- **Using 10mg per 10ml ampoule: Dilute** 25mg (25ml) with 25ml infusion fluid (i.e. total volume 50ml, 0.5mg/ml) and administer at a rate adjusted according to response
- The drug must be prepared in a syringe pump (as syringe pumps are low adsorption), and must be given via a **low-adsorption giving set** (e.g Braun Original Perfusor -Leitung PE 8723060 or Vygon Lectro-spiral 1155.80)

Bolus injection

- Only for intracoronary use
- Dose should be diluted with an equal volume of diluent

Dose in adults

Continuous intravenous infusion

- A dose of 2 to 12mg per hour is suitable although doses as high as 20mg per hour may be necessary
- Adjust according to patient's response.

Intracoronary (by bolus injection (dilute injection solution with an equal volume of diluent))

- A dose of 1mg given as a bolus injection prior to balloon inflation.
- Further doses may be given not exceeding 5mg within a 30 minute period.

Monitoring

- **Monitor** pulse and blood pressure after each rate change and during administration
- Monitor for headache, nausea and dizziness

Further information

- The drug may be given as a large volume infusion (manufacturer's preferred method) -a **low adsorption infusion bottle** (eg Braun Ecoflac or Baxter Viaflo) must be used, and it must be given via a **low adsorption giving set** (Baxter VMC9606 or Braun 8700110SP)
- However, in practice many patients are fluid restricted so this may not be practical

Storage

Store below 25°C

References

Isoket 10mg per 10ml SPC January 2023

Therapeutic classification

Nitrate drug