

Who can administer

May be administered by registered competent doctor or nurse/midwife

Important information

- **There are two intravenous ibandronic acid preparations available - Bonviva and Bondronat**
- Bonviva is licensed for the treatment of osteoporosis
- Bondronat is licensed for the treatment of Tumour Induced Hypercalcaemia, and for the prevention of skeletal related events in patients with breast cancer or bone metastases
- **Only Bonviva** is available in GUH, and this preparation should only be used for **osteoporosis**
- See under 'Dose' for adjustments required in **renal** impairment
- Ensure pre-counselling and baseline assessment for risk of adverse reactions such as **osteonecrosis of the jaw and external auditory canal**

Available preparations

Bonviva 3mg per 3mL pre-filled syringe

Bondronat 2mg per 2mL vial - not stocked in GUH

Bondronat 6mg per 6mL vial - not stocked in GUH

Reconstitution

Already in solution

Infusion fluids

Not required - product ready for use (Bonviva)

Methods of intravenous administration

Bolus intravenous injection (Bonviva only)

- Administer over 15 to 30 seconds

Dose in adults

Treatment of osteoporosis in postmenopausal women at increased risk of fracture

- Give 3mg by bolus intravenous injection every three months

Renal impairment

- Not recommended for patients with Creatinine clearance less than 30mL/minute

Further information

- All patients must receive continuous supplemental calcium and Vitamin D

- Influenza like symptoms have been reported, typically in association with the first dose
- Existing hypocalcaemia must be corrected before starting Bonviva

Storage

Store below 25°C

References

SPC 29/03/2021

Therapeutic classification

Bisphosphonate