

## Who can administer

May be administered by registered competent doctor or nurse/midwife

## Important information

### Unlicensed product

- Supplied on compassionate use by renal specialists
- Patient must be informed of unlicensed status before it is administered to them

## Available preparations

0.075mg per 1.5mL vial (0.05mg/ml)

A volume of 1.3mL is extractable from each vial

## Methods of intravenous administration

### Intravenous bolus injection

- Do not mix or dilute the injection solution prior to administration
- The drug is removed by the dialyser membrane and must be administered after blood is no longer circulating through the dialyser
- Administer by intravenous bolus injection into the venous line of the dialysis circuit at the end of each HD session
  - The dose may be given either during or after rinse back of the dialysis circuit.
  - If the dose is given **after rinse back**, administer it into the venous line followed by at least 10 mL of Sodium chloride 0.9% flush
  - If the dose is given **during rinse back**, no additional Sodium chloride 0.9% is needed to flush the line
  - The dose must be administered within 60 minutes of the completion of the syringe preparation. Discard any unused product

## Dose in adults

### Moderate-to-severe pruritus associated with chronic kidney disease in adults undergoing hemodialysis

- The recommended dose is 0.5 mcg/kg at the end of each HD session (see next point re weight to use)
- The dose/kg is determined by patient's **target dry body weight in kilograms** See Table 1.
- If a regularly scheduled HD treatment is missed, resume administration of the drug at the end of the next HD treatment

**Table 1: Injection volume based on Target Dry Weight**

Target Dry Body Weight Range (kg)	Injection volume (mL)
36-44	0.4
45-54	0.5
55-64	0.6
65-74	0.7
75-84	0.8
85-94	0.9
95-104	1
105-114	1.1
115-124	1.2
125-134	1.3
135-144	1.4
145-154	1.5
155-164	1.6
165-174	1.7
175-184	1.8
185-194	1.9
195-204	2

\* Total Injection Volume (mL) = Patient Target Dry Body Weight (kg) x 0.01, rounded to the nearest tenth (0.1 mL).

Use this formula for patient target dry body weight outside of the ranges in above table

Â

## Further information

- Difelikefalin has not been studied in patients on peritoneal dialysis and is not recommended for use in this population

## Storage

Store below 25°C

## References

Korsuva SPC 08/2021

## Therapeutic classification

Kappa opioid receptor agonist