

Who can administer

May be administered by registered competent doctor or nurse/midwife

Available preparations

Kapruvia 50micrograms per 1mL vial

Methods of intravenous administration

Intravenous bolus injection

- Do not mix or dilute the injection solution prior to administration
- The drug is removed by the dialyser membrane and must be administered after blood is no longer circulating through the dialyser
- Administer by intravenous bolus injection into the venous line of the dialysis circuit at the end of each HD session
 - The dose may be given either during or after rinse back of the dialysis circuit.
 - If the dose is given **after rinse back**, administer it into the venous line followed by at least 10 mL of Sodium chloride 0.9% flush
 - If the dose is given **during rinse back**, no additional Sodium chloride 0.9% is needed to flush the line

Dose in adults

Moderate-to-severe pruritus associated with chronic kidney disease in adults undergoing hemodialysis

- 0.5 micrograms/kg dry body weight (i.e. the target post-dialysis weight), administered at the end of each HD session
- The total dose volume (mL) required from the vial should be calculated as follows: $0.01 \times \text{dry body weight (kg)}$, rounded to the nearest tenth (0.1 mL) (see table 1 below)
- For patients with a dry body weight equal to or above 195 kg the recommended dose is 100 micrograms (2 mL)
- **If a regularly scheduled HD treatment is missed**, resume administration of the drug at the end of the next HD treatment
- **Patients with incomplete haemodialysis treatment:** for haemodialysis treatments less than 1 hour, administration of difelikefalin should be withheld until the next haemodialysis session
- **Maximum number of doses per week** is four (even if the number of dialysis sessions exceeds this)

Table 1: Injection volume based on Target Dry Weight	
Target Dry Body Weight Range (kg)	Injection volume (mL)
40-44	0.4
45-54	0.5
55-64	0.6
65-74	0.7
75-84	0.8
85-94	0.9
95-104	1
105-114	1.1
115-124	1.2
125-134	1.3
135-144	1.4
145-154	1.5
155-164	1.6
165-174	1.7
175-184	1.8
185-194	1.9
195-204	2

* Total Injection Volume (mL) = Patient Target Dry Body Weight (kg) x 0.01, rounded to the nearest tenth (0.1 mL)

Further information

- An effect of difelikefalin in reducing pruritus is expected after 2 to 3 weeks of treatment.
- Difelikefalin has not been studied in patients on peritoneal dialysis and is not recommended for use in this population

Storage

Store below 25°C

References

[Korsuva SPC](#) downloaded 15th October 2024

Therapeutic classification

Kappa opioid receptor agonist