Moxifloxacin Intravenous for Adults



Who can administer

May be administered by registered competent doctor or nurse/midwife

Important information

- Red-light antimicrobial: requires pre-authorisation by Microbiology or ID prior to use 24 hours / 7 days a week
- Consider intravenous to oral switch as soon as possible as excellent oral bioavailability (91%)
- 400mg intravenous is equivalent to 400mg orally
- There are numerous important interactions check latest BNF
- Caution (ref 1)
 - Quinolones may be associated with side effects relating to musculoskeletal, peripheral and central nervous systems, some of which may be serious, disabling and potentially permanent
 - Quinolones should be used with special caution in the elderly, patients with kidney disease, those who have had an organ transplantation or in patients being treated concomitantly with corticosteroids. These patients are at a higher risk of tendon injury
 - Patients should be informed of the risks and advised to stop treatment and contact prescriber if they
 experience pain or swelling in tendons / joints /muscle or neuropathy
- QT prolongation may occur- should not be used with other drugs that may cause QT prolongation

Available preparations

Avelox 400mg per 250ml infusion

Reconstitution

Already in solution

Infusion fluids

Not required - product ready for infusion

Methods of intravenous administration

Intermittent intravenous infusion

Administer over 60 minutes

Dose in adults

Usual dose

Give 400mg daily

Renal impairment

No dose adjustment required

Hepatic impairment

Contraindicated in impaired liver function (Child Pugh C) and in patients with transaminases increases
greater than five fold above upper limit of normal. Child-Pugh classification applies to patients
with cirrhosis. If the patient has hepatic impairment but does not have cirrhosis, dosage adjustment
should be discussed with Micro/ID

Storage

- Store below 25°C
- Do not refrigerate or freeze

References

SPC March 2021

1: AST email 25/01/2019

Therapeutic classification

Quinolone antimicrobial