

Atracurium Intravenous Infusion for Ventilated Adult Patients

Who can administer

Administration RESTRICTED - see [Appendix 1](#)

Important information

- To be used only in a critical care setting for ventilated patients
- Consideration should be given to dosing obese patients based on Ideal body weight, or an adjusted body weight (ref 1)
- As this is a neuromuscular blocker particular care and attention should be focused on the safe storage and distribution of this drug so as to minimise the risk of right patient **wrong drug**
- **This monograph relates ONLY to the use in critical care for paralysis relating to ARDS**
- For complete information and guidance refer to the manufacturers SPC on www.medicines.ie
- For Y-site compatibility [see below](#)

Available preparations

Atracurium (Tracrium) 25mg in 2.5ml ampoules

Atracurium (Tracrium) 250mg in 25ml ampoules

Reconstitution

Already in solution

Draw up using a 5 micron filter needle

Infusion fluids

Sodium chloride 0.9% or Glucose 5% (if required)

Methods of intravenous administration

Bolus intravenous injection

- Administer by rapid intravenous injection
- Administer more slowly over 1 to 2 minutes for patients with significant cardiovascular disease or hypovolaemia (ref 2)

Continuous Intravenous Infusion

- Use undiluted - 500mg in 50ml via a syringe pump

Dose in adults

Intensive care unit paralysis (eg, use for up to 48 hours in patients with early ARDS with PaO₂/FiO₂ <150, to facilitate mechanical ventilation, or for shivering from therapeutic hypothermia) (unlicensed) (ref 1)

- Initial bolus of 0.4 to 0.5mg/kg, followed by continuous IV infusion of 4 to 20 mcg/kg/**minute** (0.24 to 1.2 mg/kg/**hour**)
- Note: maximum licensed dose is 0.6mg/kg/hour
- See table below for quick reference

Â	40kg	50kg	60kg	70kg	80kg	90kg	100kg	110kg	120kg
Loading dose 0.45mg/kg	Â 18mg	22.5mg	27mgÂ	31.5mgÂ	36mgÂ	40.5mgÂ	45mgÂ	49.5mgÂ	54mgÂ
Maintenance dose	Using undiluted injection solution 500mg in 50ml- rate in ml/hour								
Â 4mcg/kg/min (0.24mg/kg/hour)	0.96	1.2	1.4	1.7Â	1.9Â	Â 2.2	2.4	Â 2.6	2.9
Â 6.6mcg/kg/min (0.4mg/kg/hour)	1.6	2	2.4Â	2.8	3.2	Â 3.6	4	Â 4.4	4.8
Â 10mcg/kg/min (0.6mg/kg/hour)	2.4	3	3.6Â	4.2	4.8	Â 5.4	6	Â 6.6	7.2Â
Â 20mcg/kg/min (1.2mg/kg/hour) (unlicensed dose once over 0.6mg/kg/hour)	4.8	6Â	7.2	8.4Â	9.6	10.8	12	Â 13.2	14.4

Storage

Store in a refrigerator at 2-8°C

References

1. Uptodate- accessed online 11 Nov 2021
2. Medusa NHS Injectable medicines accessed online 11 Nov 2021

Therapeutic classification

Neuromuscular blocking agent

BNF

Anaesthesia