

## Who can administer

Administration RESTRICTED - see [Appendix 1A](#)

## Important information

- There are numerous important interactions (eg QTc, Parkinson's) - check BNF

## Available preparations

Serenace 5mg per 1ml ampoule

## Reconstitution

Already in solution

**Draw up using a 5 micron filter needle**

## Infusion fluids

Not required - product ready for use

## Methods of intravenous administration

### **Slow intravenous injection** (ref 1)

- Administer over 3 to 5 minutes
- Maximum rate of 5mg/min in an emergency
- Can dilute with sodium chloride 0.9% or glucose 5% to produce a suitable volume to aid slow administration

## Dose in adults

### **Delirium Management in Critical Care** (ref 2)

- **Usual dose:** 2.5mg every 6 hours, adjusted according to response
- **Maximum dose:** 20mg in 24 hours
- **Older patients:** maximum of 5mg in 24 hours

## Monitoring

- Monitor blood pressure, heart rate, ECG (ref 1)
- Monitor infusion site frequently - extravasation may cause local tissue damage (ref 1)

## Storage

Store below 25°C

# References

SPC March 2011

1. Injectable Medicines administration guide, downloaded from Medusa 10/02/2025
2. UHG Critical Care Department Delirium Management algorithm V2.0 2024Â

## Therapeutic classification

Antipsychotic